Additional Information

Can the patient take any steps to protect against radiation?

If you have had a Nuclear Medicine scan in the recent past, inform the staff member to ensure that tests are not duplicated unnecessarily.

The following precautions need to be observed for 3 days after your MIBG scan:

Patients should drink plenty of fluids and empty their bladders frequently. This helps to clear the injected material from the body. Patients should observe careful hygiene when going to the toilet during this period. It is advisable to flush the toilet twice after use.

It is recommended that patients avoid spending long periods of time (in excess of 30 mins) in close contact with young children or pregnant women.

If you are a nursing mother express and discard the breast milk for 3 days after your injection. You may resume normal breast-feeding after this time.

Further Information

Directions- Beaumont Hospital:

The Nuclear Medicine Department is located on the Lower Ground Floor of Beaumont Hospital. Please check in at the Main X-ray Reception on arrival.

Who do I contact if I am unable to attend/ have queries?

If you have any questions, require further information, have specific requirements, please contact the Nuclear Medicine Secretaries on: **01-8092786.**

Young children and pregnant women should not accompany patients to Nuclear medicine.

If there is any chance you may be pregnant, please inform the Nuclear Medicine Department prior to your appointment on 01 809 2792



NMMIBG Scan Date Approved: June 2023 Version No. 2 Date for Review: June 2025 NM Department, Beaumont Hospital http://www.beaumont.ie/nm

Beaumont Imaging & Interventional Directorate





PATIENT INFORMATION

LEAFLET:

NUCLEAR MEDICINE

MIBG SCAN



Introduction

A Nuclear Medicine MIBG Scan has been requested for you and this leaflet explains what the scan involves and what you need to do to prepare for your scan. Your Nuclear Medicine scan will help make a diagnosis or monitor your treatment.

What is Radiation?

Radiation is a form of energy transmitted through space in the form of waves such as Electromagnetic radiation, eg. heat waves, visible light, radio waves or microwaves.

What is a Nuclear Medicine MIBG Scan?

This is an investigation to check for changes in the adrenal glands and surrounding tissues in the abdomen. In the Nuclear Medicine Department you will be injected with a small amount of a radioactive 'tracer' or isotope which travels to the specific part of the body requiring examination. The isotope takes approximately 4 hours to accumulate in the body so you will not be scanned until approximately 4 hours after your injection. During this waiting period you will be free to leave the hospital if you wish.

What is a gamma camera?

A gamma camera detects radiation coming from your body and forms a picture using sophisticated computer programmes. The camera consists of a flat detector which passes over the body.

Do I need to prepare for the scan?

We will send you some medication to take before the isotope injection. One tablet is to be taken 24 hours before your scan and the other is to be taken one hour before your scan. This medication protects your thyroid gland from the isotope injection.

What will happen during the scan?

After your injection you will be asked to return approximately 4 hours later for your scan.

A second scan will take place 24 hours after the injection (for example the next morning). You will be required to return to the Nuclear Medicine Department for this.

You may be asked to undress and put on a hospital gown for your scan. You will be asked to remove metal objects such as keys, coins, belt from your person prior to the scan. Your scan will be done lying down on the gamma camera.

How will the injection affect me?

You should not feel any different after your injection. The procedure will not affect your ability to drive and you do not need someone to accompany you unless you so wish.

How does a Nuclear Medicine scan differ from an X-ray?

Nuclear Medicine patients are administered substances that emit radiation and which enable staff to investigate processes that are happening in the patient's body. The amount of radiation is similar to that used in X-ray. In Nuclear Medicine the patient is emitting radiation for a period after the substance is administered. For MIBG scans, the radiation can remain in the body for up to 24 hours. Your Radiographer will advise you on the precautions you will need to apply after your scan and should be followed for 24 hours after your scan.

Will I receive much radiation?

No- the amount of radiation you receive is very small and the risks are low. The radiation dose you receive from this scan is similar to that which you would receive from your environment over a period of 1 to 2 years. Your doctor will have weighed the benefit of improved diagnosis of your condition against the small risk of the radiation exposure.